UNCLASSIFIED

Force Health Protection Brief

Indonesia

APR 2010

FORCE HEALTH PROTECTION

Indonesia

Historical diseases present in area: Diarrheal diseases, Typhoid, Hepatitis A, Malaria, Dengue Fever, Tuberculosis (TB), Chikungunya Fever, Japanese Encephalitis (JE), Shistosomiasis, and Rabies.

Recent Disease Reports: Travelers' Diarrhea**,** Avian Influenza H5N1, HIV, Malaria (*Plasmodium falciparum* 60% / *vivax* 40%), and Dengue Fever.

Force Health Protection Requirements: Up-to-date in immunizations, Pre/Post Deployment PPD for >30 day stay, anti-malarial protocols (when in rural areas), insect precautions, use approved food and water sources.

Medical infrastructure: Hospitals are not equivalent to Western standards. For severe illness or injuries utilize Commander 7th Fleet Annex "Q" requirements for medical treatment.

Environmental Hazards: Avoid congested areas due to TB and avoid local seafood (grouper, snapper, amberjack, and barracuda) as Ciguatera Fish Poisoning may be present in undercooked and well-cooked fish.

Assessment: Moderate health risk to operational forces.

26 Apr 2010

FORCE HEALTH PROTECTION

POINTS OF CONTACT

- **International SOS/TRICARE Operations Coordinator**
  - **Tel: +65-6338-9277 (Singapore)**
  - **Email: tricarelon@internationalsos.com**

- **TRICARE Area Office - Pacific**
  - **COMM: 011-81-6117-43-2036**
  - **Email: james.marron@med.navy.mil**
  - **Website: www.tricare.mil/contactus/**

- **United States Naval Forces Seventh Fleet Surgeon Liaison Officer**
  - **COMM: 808-653-2121**
  - **DSN: 315-453-2121**
  - **Email: leticia.macias@c7f.navy.mil**

FORCE HEALTH PROTECTION

Food & Water

Avoid:

Food stands

Fruits & veggies (unless peeled/cooke

Raw meat or fish

Drinking water other than bottled

Local lakes/ponds

FORCE HEALTH PROTECTION

Personal Hygiene

Frequency: 5 times/day

Timing:

- **BEFORE eating**
- **BEFORE smoking**
- **AFTER bathroom**

FORCE HEALTH PROTECTION

Avoiding Insect Bites

- **Cover up**
  - **Long sleeves/pant**
- **Insect repellent**
  - **Permethrin**
  - **DEET**
- **Sleeping**
  - **Keep bugs OUT**
  - **Bed nets, window screens**

FORCE HEALTH PROTECTION

Avian Influenza

- **Causative Agent:** Type “A” strains of influenza (flu) virus.
- **Transmission:** Contact with sick/dead poultry or contaminated surfaces.
- **Common Symptoms:** Fever, cough, sore throat, muscle aches, pneumonia, severe respiratory diseases, and other life-threatening complications.
- **Treatment:** Prescription antivirals (i.e., Tamilflu) should be effective for avian flu.
- **Prevention:** Wash hands before and after handling raw poultry and eggs, and cook poultry to 165°F.



FORCE HEALTH PROTECTION

Malaria

- **Agent –** parasite, infects the blood cells.
- **Transmission -** bite of an infected female mosquito.
- **Signs/Symptoms -** Cycles of chills, fever, sweating, vomiting and yellowing (jaundice) of the skin/eyes.
- **Prevention –** Insect skin repellent (DEET), antimalarial medications, permethrin treated bed nets/clothing and use of insecticides.



FORCE HEALTH PROTECTION

Dengue and Dengue Hemorrhagic Fever

- **Causative Agent:** Virus (*Flavivirus*).
- **Transmission:** Bites of *Aedes* mosquitoes which are day biters.
- **Symptoms:** Fever, severe headache, backache, joint pains, nausea, vomiting, eye pain and rash.
- **Treatment:** Symptomatic treatment.
- **Prevention:** No vaccine available. Use insect skin repellent (DEET), permethrin treated bed nets/clothing, and use insecticides.

Vector-borne Disease: Countermeasures

INSECT REPELLENTS FOR SKIN AND CLOTHING

DEET lotion

NSN 6840-01-284-3982

- Apply a thin coat to EXPOSED skin
- One application lasts up to 12 hours

Permethrin

- Individual Dynamic Absorption Kit (IDA)
- Treatment lasts for for over 50 launderings

NSN 6840-01-345-0237

NSN 6840-01-278-1336

- Aerosol spray can
- Treatment lasts through 5-6 washes

PERSONAL PROTECTIVE MEASURES

Permethrin On Uniform + DEET On Exposed Skin + Properly Worn Uniform = MAXIMUM PROTECTION

DOD Insect Repellent System

YOU NEED TO KNOW...
Dry cleaning removes permethrin from the uniform

FORCE HEALTH PROTECTION

Travelers' Diarrhea

- **Transmission:** Various; tainted food and water, poor sanitation.
- **Signs/Symptoms:** Abdominal cramping/pain, fever, and loose watery stools lasting 3-4 days.
- **Prevention:** Avoid local water, no raw or undercooked food, peel all fruits/veggies, avoid street vendors, and wash hands with soap prior to eating.
- **Treatment:** Antibiotics for bacterial infection (see health care provider) and Imodium AD (Loperamide). Do not take Imodium if you have blood in your stool.

FORCE HEALTH PROTECTION

Ciguatera Fish Poisoning

- **Causative Agent:** toxins produced by marine microalgae (*Gambierdiscus toxicus).*
- **Transmission:** eating fish contaminated with the toxin (barracuda, black grouper, blackfin snapper, cubera snapper, dog snapper, greater amberjack, hogfish, horse-eye jack, king mackerel, and yellowfin grouper have been known to carry ciguatoxins).
- **Symptoms:** nausea, vomiting, and neurologic symptoms such as tingling fingers or toes. Cold things may feel hot and hot things may feel cold.
- **Treatment:** symptomatic treatment. Ciguatera has no cure. Symptoms usually go away in days or weeks but can last for years.
- **Prevention:** avoid local seafood as Ciguatera Fish Poisoning may be present in undercooked and well-cooked fish.

FORCE HEALTH PROTECTION

Tuberculosis (TB)

- **Causative Agent:** Bacteria (*Mycobacterium tuberculosis*).
- **Transmission:** Airborne droplet via respiration.
- **Symptoms:** Cough, fatigue, night sweats, fever and rapid heart rate.
- **Treatment:** Antibiotics (some strains resistant to multiple drugs) and respiratory isolation.
- **Prevention:** No vaccine available. Avoid contact with persons with active TB, congested areas, and receive annual screening (PPD).



M. tuberculosis

FORCE HEALTH PROTECTION

Sexually Transmitted Disease

HIGH RISK FOR STDs - Hepatitis B, HIV, etc.

FHP Requirements

- **Avoid sexual contact.**
- **Use latex condoms if sexually active.**
- **Use Condom Sense!**

***STD diseases can be transmitted by tattoo needles.**

FORCE HEALTH PROTECTION

Local Animals

- **Many countries have NO rabies control programs!**
- **Do not feed or attempt to touch animals.**
- **Report bites immediately.**
- **No mascots!**

VENOMOUS SNAKES

COBRAS/KRAITS/CORAL SNAKES

Short fangs fixed in erect position in front of mouth.

Round eye pupils.



Head small and not distinct from the body.

PIT

Long, hinged fangs that tuck into roof of mouth when not in use.



Vertically elliptical eye pupils; pit vipers with heat sensing organ between eye and nostril.



Broad triangular head very distinct from narrow neck.



HAZARDOUS PLANTS

- Plant resins cause skin irritation
- Poisonous roots, stems, leaves, and fruit
- Weeds and stinging nettles
- Thorny shrubs and trees

Poisonous Plants

Mucuna pruritum

Systemic

Thorny Plant

Punctures and Abrasions

COUNTERMEASURES

- Avoid touching unfamiliar plants
- Use clothing as protective barrier
- Rinse skin promptly after exposure
- Wash clothing after contact
- Never eat any part of unfamiliar

Required Health Measures

- Pages 13-15 USNS Mercy Handbook
- Immunizations
- Malaria chemoprophylaxis
  - Terminal chemoprophylaxis*
- Mosquito & insect bite prevention
- Pre-mission health assessment
  - Include Tuberculosis (TB) screening test
  - G6PD blood test*
- Obtain personal medication

*Depends on itinerary – obtain specific guidance

Required Immunizations

- Start vaccinations early to accommodate vaccines that require several doses within a series
- Mission-specific:
  - Hepatitis B*/**
  - Typhoid
  - Japanese Encephalitis**
  - Rabies**
- Routine:
  - Tetanus/diphtheria
  - Measles (MMR)
  - Polio
  - Varicella (chicken pox)
  - H1N1 Influenza
  - Seasonal Influenza
  - Hepatitis A

* Required for health care providers

** For high risk groups only – obtain specific guidance

FORCE HEALTH PROTECTION

Indonesia Entry Requirements

- **Although yellow fever is not a disease risk in Indonesia, the government requires travelers arriving from countries where Yellow Fever is present (see guidance for countries in Africa, Central and South America) to present proof of yellow fever vaccination.**

- **Example of Yellow Fever card.**



sceau du Service de la Santé Publique des États Unis, ou le timbre spécial "S-C" approuvé par ce service.)
Toute correction ou rature sur le certificat ou l'omission d'une quelconque des mentions qu'il comporte peut affecter sa validité.

*If unsuccessful, vaccination must be repeated and a new certificate executed.
Si la vaccination n'a pas prise, il faudra recommencer et un nouveau certificat devra être établi.

INTERNATIONAL CERTIFICATE OF VACCINATION OR REVACCINATION AGAINST YELLOW FEVER

CERTIFICAT INTERNATIONAL DE VACCINATION OU DE REVACCINATION CONTRE LA FIÈVRE JAUNE

This is to certify that
Je soussigné(e) certifie que ..... sex / sexe .....

whose signature follows
dont la signature suit ..... date of birth / né(e) le .....

has on the date indicated been vaccinated or revaccinated against yellow fever.
a été vacciné(e) ou revacciné(e) contre la fièvre jaune à la date indiquée.

Date	Signature and professional status of vaccinator Signature et qualité professionnelle du vaccinateur	Origin and batch number of vaccine Origine du vaccin employé et numéro du lot	Official stamp of vaccinating center Cachet officiel du centre de vaccination
1/17/61	Capt. John P. Leddy, MC Walter Reed Army Medical Center, Washington, D.C.	National Drug Co. lot # 4798	MEDICAL SERVICES
6 Jun 62	Alvan Burgess Capt MC	National Dy #5319	

VACCINATING CENTER
CENTRE DE VACCINATION

ADDRESS / ADRESSE (CITY—VILLE) (STATE—ÉTAT)

THIS CERTIFICATE IS VALID only if the vaccine used has been approved by the World Health Organization and if the vaccinating center has been designated by the health administration for the country in which that center is situated.

THE VALIDITY OF THIS CERTIFICATE shall extend for a period of 6 years, beginning 10 days after the date of vaccination **(for India, Pakistan, and Ceylon 12 days)** or, in the event of a revaccination, within such period of 6 years, from the date of that revaccination.

Any amendment of this certificate, or erasure, or failure to complete any part of it, may render it invalid.

CE CERTIFICAT N'EST VALABLE que si le vaccin employé a été approuvé par l'Organisation Mondiale de la Santé et si le centre de vaccination a été habilité par l'administration sanitaire du territoire dans lequel ce centre est situé.

LA VALIDITÉ DE CE CERTIFICAT couvre une période de six ans commençant dix jours après la date de la vaccination (pour l'Inde, Pakistan et Ceylan 12 jours) ou, dans le cas d'une revaccination au cours de cette période de six ans, le jour de cette revaccination.

Toute correction ou rature sur le certificat ou l'omission d'une quelconque des mentions qu'il comporte peut affecter sa validité.

U. S. GOVERNMENT PRINTING OFFICE : 1956—O-404968

Questions?

- **Please contact your local health care provider**

- **Navy Environmental and Preventative Medicine Unit No. 6**

  Duty Phone: 808-471-0237

  nepmu6admin@med.navy.mil

- **Navy Environmental and Preventative Medicine Unit No. 5**

  Duty Phone: 619-556-7070

  nepmu5@med.navy.mil